

COSMOLOGY WITH QUANTUM MATTER AND A CLASSICAL GRAVITATIONAL FIELD

Marcel Reginatto

*Physikalisch-Technische Bundesanstalt,
Bundesallee 100, 38116 Braunschweig, Germany*

(Dated: August 5, 2009)

Abstract

I consider a hybrid cosmological model with a classical gravitational field and a quantized massive scalar field in spherically symmetric gravity. The interaction between classical and quantum fields is described using the formalism of ensembles on configuration space, an approach that provides an alternative to semiclassical gravity. After giving an outline of the formulation as it applies to spherically symmetric gravity, I discuss a particular solution that is analogous to the Einstein universe of classical relativity.

PACS numbers: 04.50.+h, 04.62.+v

I. INTRODUCTION

There are a few good reasons to consider hybrid systems in which the gravitational field remains classical while matter is assumed to consist of quantum fields. A full theory of quantum gravity is not yet available, and an approximation in which spacetime remains classical while matter is described in terms of quantum fields is often physically and computationally appropriate. Furthermore, since the quantization of gravity does not appear to follow from consistency arguments alone [1], it is of interest to investigate to what extent a hybrid system may provide a consistent, satisfactory description of matter and gravitation. The study of such systems can provide valuable clues that may help in the search for a full quantum theory of gravity. Finally, one must also consider the possibility that the gravitational field may not be quantum in nature [2, 3, 4]. For example, Butterfield and Isham, while putting forward the point of view that some type of theory of quantum gravity should be sought, have concluded that there is arguably no definitive proof that general relativity has to be quantized [5]. Dyson has argued that it might be impossible in principle to observe the existence of individual gravitons, and this has lead him to the conjecture that “the gravitational field described by Einstein’s theory of general relativity is a purely classical field without any quantum behaviour” [6]. His observations regarding the impossibility of detecting gravitons have been confirmed recently by detailed calculations [7, 8]. If Dyson’s conjecture is true, hybrid models become unavoidable.

In the standard approach used for coupling quantum fields to a classical gravitational field (i.e., semiclassical gravity), the energy momentum tensor that serves as the source in the Einstein equations is replaced by the expectation value of the energy momentum operator $\hat{T}_{\mu\nu}$ with respect to some quantum state Ψ :

$${}^4R_{\mu\nu} - \frac{1}{2}g_{\mu\nu} {}^4R + \lambda g_{\mu\nu} = \frac{\kappa}{2} \langle \Psi | \hat{T}_{\mu\nu} | \Psi \rangle$$

where ${}^4R_{\mu\nu}$ is the curvature tensor, 4R the curvature scalar and $g_{\mu\nu}$ the metric tensor in spacetime, λ is the cosmological constant and $\kappa = 16\pi G$ (in units where $c = 1$) where G is the gravitational constant. This approach, however, presents a number of well known difficulties. An alternative to semiclassical gravity is provided by the formalism of ensembles on configuration space [9]. The formalism can be applied to both quantum and classical systems and allows a general and consistent description of interactions between them. When applied to standard quantum mechanics, it can be shown that the approach overcomes difficulties arising in previous attempts; in particular, the correct equations of motion for the quantum and classical sectors are recovered in the limit of no interaction, conservation of probability and energy are satisfied, uncertainty relations hold for conjugate quantum variables, and the formalism allows a back reaction of the quantum system on the classical system [10, 11]. The approach can be generalized to describe the coupling of a quantized field to a classical gravitational field; it is therefore an appropriate formalism to use if one wants to investigate hybrid cosmological models.

In the next section, I describe the coupling of a quantized scalar field to a classical gravitational field using ensembles on configuration space. I then consider midisuperspace

cosmological models in spherically symmetric gravity and derive a particular solution which describes a static universe. In the last section, I summarize the results and give conclusions.

II. EQUATIONS FOR A QUANTIZED SCALAR FIELD INTERACTING WITH A CLASSICAL METRIC FIELD

In the formalism of ensembles on configuration, the state of a system is described in terms of a probability density P together with its canonically conjugate variable S and the equations of motion are derived from an ensemble Hamiltonian. I first illustrate the approach for the case of vacuum gravity [12, 13, 14] and then discuss the formulation for hybrid systems. The most direct way of introducing a classical configuration space ensemble for gravity is to start from the Einstein-Hamilton-Jacobi equation, which in the metric representation takes the form [15]

$$\mathcal{H}_h = \kappa G_{ijkl} \frac{\delta S}{\delta h_{ij}} \frac{\delta S}{\delta h_{kl}} - \frac{1}{\kappa} \sqrt{h} (R - 2\lambda) = 0, \quad (1)$$

where R is the curvature scalar and h_{kl} the metric tensor on a three-dimensional spatial hypersurface, and $G_{ijkl} = (2h)^{-1/2} (h_{ik}h_{jl} + h_{il}h_{jk} - h_{ij}h_{kl})$ is the DeWitt supermetric. The functional S is assumed to be invariant under the gauge group of spatial coordinate transformations, which is equivalent to satisfying the momentum constraints of the canonical formulation of general relativity. Eq. (1) corresponds to an infinity of constraints, one at each point. It is possible to introduce an alternative viewpoint [16, 17] in which Eq. (1) is regarded as an equation to be integrated with respect to a “test function” in which case we are dealing with one equation for each choice of lapse function N ,

$$\int d^3x \, N \mathcal{H}_h = 0; \quad (2)$$

i.e. for each choice of foliation. Such an alternative viewpoint is extremely useful: although it may be impossible to find the general solution (which requires solving the Einstein-Hamilton-Jacobi equation for all choices of lapse functions), it may be possible to find particular solutions for specific choices; for example, the choice $S \sim \int d^3x \sqrt{h}$ is a particular solution that describes de Sitter spacetime in a flat foliation [17].

An appropriate *ensemble* Hamiltonian for vacuum gravity is given by

$$\tilde{H}_h = \int d^3x \int Dh \, P \, N \mathcal{H}_h \quad (3)$$

(technical issues are discussed in more detail in [14]). The functional P is also assumed to be invariant under the gauge group of spatial coordinate transformations. The corresponding equations have the form

$$\frac{\partial P}{\partial t} = \frac{\delta \tilde{H}_h}{\delta S}, \quad \frac{\partial S}{\partial t} = -\frac{\delta \tilde{H}_h}{\delta P}$$

where $\delta/\delta F$ denotes the variational derivative with respect to the functional F [12]. Assuming the constraints $\frac{\partial S}{\partial t} = \frac{\partial P}{\partial t} = 0$, these equations lead to Eq. (2), as required, and to a

continuity equation,

$$\int d^3x N \frac{\delta}{\delta h_{ij}} \left(P G_{ijkl} \frac{\delta S}{\delta h_{kl}} \right) = 0. \quad (4)$$

These two equations define an ensemble on configuration space for the case of vacuum gravity.

A hybrid system where a quantum scalar field ϕ couples to the classical metric h_{kl} requires a generalization of Eq. (3) in which [9]

$$\tilde{H}_{\phi h} = \int d^3x \int Dh P N [\mathcal{H}_{\phi h} + F_\phi], \quad (5)$$

where

$$\mathcal{H}_{\phi h} = \mathcal{H}_h + \frac{1}{2\sqrt{h}} \left(\frac{\delta S}{\delta \phi} \right)^2 + \sqrt{h} \left[\frac{1}{2} h^{ij} \frac{\partial \phi}{\partial x^i} \frac{\partial \phi}{\partial x^j} + V(\phi) \right]$$

is the ensemble Hamiltonian for gravity with a classical scalar field and

$$F_\phi = \frac{\hbar^2}{4} \frac{1}{2\sqrt{h}} \left(\frac{\delta \log P}{\delta \phi} \right)^2$$

is an additional, non-classical kinetic energy term. Assuming again the constraints $\frac{\partial S}{\partial t} = \frac{\partial P}{\partial t} = 0$, the corresponding equations are given by

$$\int d^3x N \left[\mathcal{H}_{\phi h} - \frac{\hbar^2}{2\sqrt{h}} \left(\frac{1}{A} \frac{\delta^2 A}{\delta \phi^2} \right) \right] = 0,$$

where $A \equiv \sqrt{P}$, and a continuity equation that is identical to Eq. (4).

III. HYBRID COSMOLOGICAL MODEL IN SPHERICALLY SYMMETRIC GRAVITY

A. Equations

I consider a midisuperspace hybrid cosmological model in spherically symmetric gravity. In the case of spherical symmetry, the line element may be written in the form

$$g_{\mu\nu} dx^\mu dx^\nu = -N^2 dt^2 + \Lambda^2 (dr + N^r dt)^2 + R^2 d\Omega^2. \quad (6)$$

The lapse function N and the shift function N^r are functions of the radial coordinate r and the time coordinate t . The configuration space for the gravitational field consists of two fields, R and Λ . Under transformations of r , R behaves as a scalar and Λ as a scalar density. Spherically symmetric gravity is discussed in detail in a number of papers, mostly in reference to the canonical quantization of black hole spacetimes. For discussions using the metric representation, see for example [18, 19, 20]. For discussions of the Einstein-Hamilton-Jacobi equation in the context of the WKB approximation of quantized spherically symmetric gravity, see for example [21, 22].

I now set $\hbar = c = G = 1$. The Einstein-Hamilton-Jacobi equation for the case of a vacuum gravity takes the form

$$\mathcal{H}_{\Lambda R} = -\frac{1}{R} \frac{\delta S}{\delta R} \frac{\delta S}{\delta \Lambda} + \frac{1\Lambda}{2R^2} \left(\frac{\delta S}{\delta \Lambda} \right)^2 + \lambda \frac{\Lambda R^2}{2} + V = 0$$

where S is assumed to be invariant under diffeomorphisms. V is related to the curvature scalar by ${}^4R = -4\Lambda R^2 V$, and is given by

$$V = \frac{RR''}{\Lambda} - \frac{RR'\Lambda'}{\Lambda^2} + \frac{R'^2}{2\Lambda} - \frac{\Lambda}{2} \quad (7)$$

where primes indicate derivatives with respect to r .

The ensemble Hamiltonian of a hybrid system where the matter is in the form of a minimally coupled quantized radially symmetric scalar field of mass m is given by

$$\tilde{H}_{\phi\Lambda R} = \int dr \int Dh P N [\mathcal{H}_{\phi\Lambda R} + F_\phi], \quad (8)$$

where

$$\mathcal{H}_{\phi\Lambda R} = \mathcal{H}_{\Lambda R} + \frac{1}{2\Lambda R^2} \left(\frac{\delta S}{\delta \phi} \right)^2 + \frac{R^2}{2\Lambda} \phi'^2 + \frac{\Lambda R^2 m^2}{2} \phi^2,$$

and

$$F_\phi = \frac{1}{8\Lambda R^2} \left(\frac{\delta \log P}{\delta \phi} \right)^2.$$

Eq. (8) is the analogous of Eq. (5) for the case of spherically symmetric gravity. Assuming again the constraints $\frac{\partial S}{\partial t} = \frac{\partial P}{\partial t} = 0$, the corresponding equations are

$$\int dr N \left[\mathcal{H}_{\phi\Lambda R} - \frac{1}{2\Lambda R^2} \left(\frac{1}{A} \frac{\delta^2 A}{\delta \phi^2} \right) \right] = 0, \quad (9)$$

where $A \equiv P^{1/2}$, and the continuity equation

$$\int dr N \left[\frac{\delta}{\delta R} \left(P \frac{1}{R} \frac{\delta S}{\delta \Lambda} \right) + \frac{\delta}{\delta \Lambda} \left(P \frac{1}{R} \frac{\delta S}{\delta R} - P \frac{\Lambda}{R^2} \frac{\delta S}{\delta \Lambda} \right) - \frac{\delta}{\delta \phi} \left(P \frac{1}{\Lambda R^2} \frac{\delta S}{\delta \phi} \right) \right] = 0. \quad (10)$$

B. A particular solution

I now discuss a particular solution of these equations. I am interested in a hybrid cosmological model where the metric is that of a closed Friedmann universe. For computational reasons, it will be useful to consider the coordinate system given by

$$N = 1, \quad N^r = 0, \quad \Lambda = a_0, \quad R = a_0 \sin r, \quad (11)$$

where the line element is of the form of Eq. (6) with $r \in [0, 2\pi)$. This metric describes a closed Friedmann universe with scale factor a_0 . The particular solution that I will consider is one that describes a Friedmann universe in a foliation of spaces of constant positive

curvature. The advantage of such a choice of foliation is that it leads to a potential term, Eq. (7), which simplifies to

$$V = -\frac{3R^2}{2\Lambda} = -\frac{3a_0 \sin^2 r}{2}. \quad (12)$$

where the second equality is only valid for the particular coordinate conditions described by Eqs. (11).

Even with the assumption of a foliation of spaces of constant positive curvature, a solution of equations Eqs. (9) and (10) remains difficult. To solve these equations, I consider an ansatz of the form

$$S[R, \Lambda, \phi] = 0, \quad P[R, \Lambda, \phi] = P[\phi | R, \Lambda] P[R, \Lambda]$$

with

$$P^{1/2}[\phi | R, \Lambda] \equiv A_{(0)} = C \exp \left\{ -a_0^6 \int \int dy dz \sin^2 y \sin^2 z \phi_y K_{yz} \phi_z \right\}.$$

where C is a normalization factor, the kernel K_{xy} is real and symmetric, $K_{xy} = K_{yx}$, and the notation $P[A | B]$ denotes a conditional probability.

This is a very special solution: since Eq. (10) is linear in S , the choice $S = 0$ solves this equation independent of the form of A . From a physical point of view, this assumption corresponds to selecting a static solution (see the discussion in the Appendix). With this choice of S , the functional $A_{(0)}$ obeys a functional Schrödinger equation in a space of constant curvature, as shown below. The form of $A_{(0)}$ chosen here is that of a ground state Gaussian functional. Instead of $A_{(0)}$, one may also consider the excited states which solve the functional Schrödinger equation; I discuss the consequences of making this alternative choice at the end of this section.

If I calculate the second functional derivative of $A_{(0)}$, I get an expression of the form

$$\frac{1}{A_{(0)}} \frac{\delta^2 A_{(0)}}{\delta \phi_r^2} = -2a_0^6 \sin^4 r K_{rr} + 4a_0^{12} \sin^4 r \int \int dy dz \sin^2 y \sin^2 z \phi_y K_{yr} K_{rz} \phi_z. \quad (13)$$

To solve these equations, I will choose a lapse function N which is constant. Combining Eqs. (9), (12) and (13) and collecting terms that have the same powers of ϕ , I get the two equations

$$\int dr \sin^2 r \left[\left(\frac{\lambda a_0^3}{2} - \frac{3a_0}{2} \right) + a_0^3 K_{rr} \right] = 0, \quad (14)$$

and

$$\int dr \sin^2 r \left[\frac{a_0}{2} \phi'^2 + \frac{a_0^3 m^2}{2} \phi^2 - 2a_0^9 \int \int dy dz \sin^2 y \sin^2 z \phi_y K_{yr} K_{rz} \phi_z \right] = 0. \quad (15)$$

After an integration by parts in Eq. (15), the equations are of a type that is standard in the context of the Schrödinger functional representation of quantum field theory in curved spacetimes [23, 24, 25, 26]. I solve this functional Schrödinger equation in two steps.

I first consider Eq. (15). The solution of the kernel K_{xy} is given by [26]

$$K_{xy} = \frac{1}{2a_0^4} \sum_n \sqrt{\gamma_n} \psi_x^{(n)} \psi_y^{(n)},$$

where the basis functions $\psi_r^{(n)}$ are solutions of a Schrödinger-type equation in a space of constant curvature,

$$-\frac{1}{\sin^2 r} \frac{\partial}{\partial r} \left(\sin^2 r \frac{\partial \psi_r^{(n)}}{\partial r} \right) + m^2 a_0^2 \psi_r^{(n)} = \gamma_n \psi_r^{(n)}.$$

The $\psi_n(r)$ satisfy orthonormality and completeness relations. The eigenvalues γ_n are given by

$$\gamma_n = n^2 - 1 + m^2 a_0^2, \quad n = 1, 2, 3, \dots$$

C. Renormalization

Given a solution of Eq. (15), one can use Eq. (14) to express a_0 in terms of the cosmological constant and the energy E_0 of the quantized scalar field, since [26]

$$E_0 \sim a_0^3 \int dr \sin^2 r K_{rr}.$$

However, $\int dr \sin^2 r K_{rr} \sim \sum_n \gamma_n$, which diverges. This is a consequence of the infinite zero-point energy of the quantum field. It is necessary to use a renormalization procedure to extract a finite result.

In this particular case, the equation for $A_{(0)}$ is similar in form to a functional Schrödinger equation for a quantum scalar field in an Einstein universe, so it is possible to use previous results from the literature where renormalization procedures have been carried out for this problem (for a thorough analysis, see reference [27]). Note that this is not the generic case: since the equations of hybrid cosmology are not linear, in general they will not map to a Schrödinger-type functional equation. The simplification that is achieved here is a direct consequence of choosing $S = 0$ and a foliation of spaces of constant positive curvature. With a different ansatz, the equations can not be solved in this way.

A detailed study of this solution will be the subject of a future publication. In this note, I will only make a few remarks concerning some qualitative aspects. Renormalization leads to a finite expression for the energy E_0^{ren} that has the functional form [27]

$$E_0^{ren} \sim \frac{1}{a_0} \alpha(ma_0)$$

where the function α depends on the dimensionless quantity ma_0 in a complicated way. Using the renormalized value in Eq. (14) and rescaling α by a constant factor leads to

$$\lambda a_0^4 - 3a_0^2 + \alpha(ma_0) = 0.$$

In the case $m = 0$, α is constant and takes a negative value. Then, the equation has one real solution,

$$a_0^2 = \frac{1}{2\lambda} \left(3 + \sqrt{9 + 4\lambda|\alpha(0)|} \right).$$

The case $m \neq 0$ is much more complicated because α is no longer a constant. There are values of m with $\alpha > 0$, in which case it is possible to find solutions with $\lambda = 0$ and $a_0 = \sqrt{\alpha/3}$. The classical Einstein universe requires fine-tuning λ to get a static solution and it is therefore not stable. However, it is not clear whether there are stable solutions among these static hybrid cosmological models which do not require a cosmological constant.

A similar analysis may be carried out where the ground state functional $A_{(0)}$ is replaced by an excited state. Consider a first excited state $A_{(1)}$ specified by the eigenfunction $\psi_r^{(n)}$. This state will differ in energy from the (divergent) ground state energy by a finite amount $\Delta E(n)$ which depends on the value of γ_n , with $\Delta E(n) = \sqrt{\gamma_n}$ [26]. $\Delta E(n)$ can only take discrete values because γ_n is quantized, and this means that a_0 will also be restricted to discrete values. Therefore, the coupling of the quantum scalar field and classical gravitational field appears to lead to the quantization of the radius of the universe.

IV. CONCLUDING REMARKS

I give a brief summary of the main results. A description of a hybrid cosmological model consisting of a classical gravitational field and a quantized massive scalar field has been given using the formalism of ensembles on configuration space. For the case of spherically symmetric gravity, a particular static solution was found that is analogous to the Einstein universe of classical relativity.

The hybrid cosmological model examined here was previously considered in reference [9], but only within the minisuperspace approximation. One of the aims of the present work was to reexamine this model using the next level of approximation; i.e., in the context of a midisuperspace formulation like spherically symmetric gravity. The conclusion is that the space of solutions is very rich by comparison. This is a consequence of the renormalization procedure that needs to be carried out to achieve a finite result. This complexity is absent from the minisuperspace solution. A common feature of both levels of approximation is that the coupling of the quantum scalar field and classical gravitational field leads to the quantization of the radius of the universe, a remarkable result. It would be interesting to know whether this is a generic feature of hybrid closed cosmological models or whether this is a feature that is particular to the solution considered here. In the minisuperspace solution, it is clear that a singular solution with $a = 0$ is excluded; the situation is more complicated in the case of the midisuperspace solution and it appears that $a = 0$ is possible for some values of the mass m .

In the Einstein universe of classical relativity, the cosmological constant acts as a repulsive force which balances the gravitational attraction of matter; it is essential to include it in order to achieve a static solution. The Einstein universe is therefore unstable. In the minisuperspace solution considered here, it is possible to achieve a static solution with a vanishing cosmological constant for some values of the mass m and it would appear that quantum fluctuations are sufficient to balance the gravitational attraction and provide equilibrium under some circumstances. It is not clear however whether such solutions are stable under perturbation.

Finally, it is worth pointing out that there is a natural, well defined intrinsic concept

of time in this formulation that can be used to evolve the fields through a given choice of foliation, as shown in the Appendix. This is a consequence of keeping the gravitational field classical, and it leads to rate equations for the fields that agree with those derived using the Hamilton-Jacobi formulation of classical relativity.

Acknowledgments

I thank M. J. W. Hall and Stephen Boughn for valuable discussions.

APPENDIX: CONTINUITY EQUATION AND RATE EQUATIONS

The purpose of this appendix is to point out that the interpretation of the equation

$$\int d^3x N \frac{\delta}{\delta h_{ij}} \left(P G_{ijkl} \frac{\delta S}{\delta h_{kl}} \right) = 0$$

as a continuity equation leads to a rate equation that relates $\frac{\partial h_{kl}}{\partial t}$ and $\frac{\delta S}{\delta h_{kl}}$. This follows from the observation that such an interpretation is only possible if the “field velocity” $\frac{\partial h_{kl}}{\partial t}$ is related to $G_{ijkl} \frac{\delta S}{\delta h_{kl}}$ in a linear fashion. The most general rate equation for the metric h_{ij} that is consistent with such a linear relationship is of the form

$$\delta h_{ij} = \left(\gamma G_{ijkl} \frac{\delta S}{\delta h_{kl}} + \delta_\epsilon h_{ij} \right) \delta t \quad (\text{A.1})$$

where γ is an arbitrary function. I have include in Eq. (A.1) a term $\delta_\epsilon h_{kl} = -(D_k \epsilon_l + D_l \epsilon_k)$ which allows for gauge transformations of h_{kl} , which is permitted because the gauge transformations are assumed to leave $\int DhP$ invariant.

Eq. (A.1) agrees with the usual rate equations [28] provided we identify γ with the lapse function N and ϵ_k with the negative of the shift vector N_k . In particular, if the shift vector is set to zero and $\frac{\delta S}{\delta h_{kl}} = 0$, it follows that $\frac{\partial h_{ij}}{\partial t} = 0$ and we are dealing with a static metric.

A similar result holds in the case of matter fields (classical or quantum) interacting with a classical metric field. This shows that there is a natural, well defined intrinsic concept of time that can be used to evolve the fields through a given choice of foliation.

-
- [1] M. Albers, C. Kiefer and M. Reginatto, *Measurement analysis and quantum gravity*, Phys. Rev. D **78**, 064051 (2008)
 - [2] L. Rosenfeld, *On Quantization of Fields*, Nucl. Phys. **40**, 353 (1963).
 - [3] S. Carlip, *Is quantum gravity necessary?*, Class. Quantum Grav. **25** 154010 (2008).
 - [4] S. Boughn, *Nonquantum Gravity*, Found. Phys., **39**, 331 (2009)
 - [5] J. Butterfield and C. Isham, *Spacetime and the philosophical challenge of quantum gravity*. In: *Physics meets philosophy at the Planck scale*, C. Callender and N. Huggett, eds. (Cambridge University Press, 2001), Chap. 2.

- [6] F. Dyson, *The World on a String*, review of *The Fabric of the Cosmos: Space, Time, and the Texture of Reality* by Brian Greene, New York Review of Books, 51(8) (2004).
- [7] T. Rothman and S. Boughn, *Can Gravitons Be Detected?*, Found. Phys. 36, 1801 (2006).
- [8] S. Boughn and T. Rothman, *Aspects of graviton detection: graviton emission and absorption by atomic hydrogen*, Class. Quantum Grav. **23** 5839 (2006).
- [9] M. J. W. Hall and M. Reginatto, *Interacting classical and quantum ensembles*, Phys. Rev. A **72**, 062109 (2005)
- [10] M. J. W. Hall, *Consistent classical and quantum mixed dynamics*, Phys. Rev. A **78**, 042104 (2008)
- [11] M. Reginatto and M. J. W. Hall, *Quantum-classical interactions and measurement: a consistent description using statistical ensembles on configuration space*, J. Phys.: Conf. Ser. 174 012038 (2009)
- [12] M.J.W. Hall, K. Kumar and M. Reginatto, *Bosonic field equations from an exact uncertainty principle*, J. Phys. A **36**, 9779 (2003).
- [13] M. J. W. Hall, *Exact uncertainty approach in quantum mechanics and quantum gravity*, General Relativity and Gravitation **37** 1505 (2005).
- [14] M. Reginatto, *Exact uncertainty principle and quantization: implications for the gravitational field*, Braz. J. Phys. **35**, 476 (2005).
- [15] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [16] C. Kiefer, *What is the geometry of superspace?*, in from Newton's bucket to quantum gravity, ed. by J. B. Barbour and H. Pfister, Birkhuser, Boston (1993).
- [17] C. Kiefer, *The semiclassical approximation to quantum gravity*, Report Freiburg THEP-93/27 and gr-qc/9312015 (1993).
- [18] J. D. Romano, *Spherically Symmetric Scalar Field Collapse: An Example of the Spacetime Problem of Time*, arXiv:gr-qc/9501015 v1 14 Jan 1995.
- [19] K. Kuchař, *Geometrodynamics of Schwarzschild black holes*, Phys. Rev. **D50**, (1994) 3961.
- [20] S. R. Lau, *On the Canonical Reduction of Spherically Symmetric Gravity*, gr-qc/9508028 v3 14 Feb 1996
- [21] W. Fischler, D. Morgan and J. Polchinski, *Quantization of false-vacuum bubbles: A Hamiltonian treatment of gravitational tunneling*, Phys. Rev. **D42** (1990) 4042.
- [22] T. Brotz and C. Kiefer, *Semiclassical black hole states and entropy*, Phys. Rev. D 55, 2186-2191 (1997).
- [23] C. Kiefer, *Quantum Gravity* (Clarendon Press, Oxford, 2004)
- [24] R. Jackiw, *Analysis On Infinite Dimensional Manifolds – Schrödinger Representation For Quantized Fields,* in *Field Theory and Particle Physics* (O.Eboli, et al. eds.). World Scientific, Singapore 1990. Reprinted in R. Jackiw, *Diverse Topics in Theoretical and Mathematical Physics*, pp 383-445. World Scientific, Singapore 1995.
- [25] B. Hatfield, *Quantum field theory of point particles and strings*, Perseus Books, Cambridge, Massachusetts (1992).
- [26] D. V. Long and G. M. Shore, *The Schrödinger wave functional and vacuum states in curved spacetime*, Nucl. Phys. B **530**, 247 (1998).
- [27] C. A. R. Herdeiro, R. H. Ribeiro and M. Sampaio, *Scalar Casimir effect on a D-dimensional*

- Einstein static universe*, Class. Quantum Grav. **25**, 165010 (2008).
- [28] U. H. Gerlach, *Derivation of the Ten Einstein Field Equations from the Semiclassical Approximation to Quantum Geometrodynamics*, Phys. Rev. 177, 1929 (1969)